

ON THE SET OF t -LINKED OVERRINGS OF AN INTEGRAL DOMAIN

A. MIMOUNI

ABSTRACT. Let R be an integral domain with quotient field L . An overring T of R is t -linked over R if $I^{-1} = R$ implies that $(T : IT) = T$ for each finitely generated ideal I of R . Let $O_t(R)$ denotes the set of all t -linked overrings of R and $O(R)$ the set of all overrings of R . The purpose of this paper is to study some finiteness conditions on the set $O_t(R)$. Particularly, we prove that if $O_t(R)$ is finite, then so is $O(R)$ and $O_t(R) = O(R)$, and if each chain of t -linked overrings of R is finite, then each chain of overrings of R is finite. This yields that the t -linked approach is more efficient than the Gilmer's treatment in [23]. We also examine the finiteness conditions in some Noetherian-like settings such as Mori domain, quasicohherent Mori domain, Krull domain etc. We establish a connection between $O_t(R)$ and the set of all strongly divisorial ideals of R and we conclude by a characterization of domains R that are t -linked under all their overrings.

1. INTRODUCTION

Throughout R is an integral domain (which is not a field) with quotient field L . By an overring of R we mean an integral domain T such that $R \subseteq T \subseteq L$. In their study of the residually algebraic pairs of integral domains in [3], Ayache and Jaballah encountered the following two conditions on the set of overrings of an integral domain R :

- (i) R has only finitely many overrings.
- (ii) Each chain of distinct overrings of R is finite.

In extending results of [3] in [25] and [26], Jaballah asked [26, Question 1] for a characterizations for domains with condition (i), that is, domains with finitely many overrings. In [23], R. Gilmer labels the above two conditions as (FO) and (FC) in the following meaning:

- (i) R is an FO -domain if R has finitely many overrings;
- (ii) R is an FC -domain if each chain of distinct overrings of R is finite.

He completely characterizes these domains [23, Theorem 2.14, and Theorem 3.4]. Gilmer's characterizations involve the integral closure R' of R , the conductor $(R : R')$ and the notion of FIP property (stands for "finitely many intermediate R -algebras") introduced and investigated by D. Anderson, D. Dobbs and B. Mullins in [1]. In a recent work, A. Jaballah gives an algorithm of how to compute the number of overrings of an integrally closed domain with finitely many overrings

2000 *Mathematics Subject Classification.* Primary 13G05, 13F05; Secondary 13B02, 13B22.

Key words and phrases. t -linked overring, finite chain of t -linked overrings, FO -domain, FC -domain, Prüfer domain, Mori domain, strongly divisorial ideal.

This work was funded by KFUPM under Project # FT/18-2005.

[27], however the question is still open for an arbitrary integral domain.

According to [11], an overring T of R is said to be t -linked over R if $I^{-1} = R$ for a finitely generated ideal I of R implies that $(IT)^{-1} = (T : IT) = T$. The t -linked concept was used in [11, Theorem 2.10, and Corollary 2.18] to find characterizations of certain classes of $PVMD$'s analogous to characterizations of Prüfer domains due to Davis and Richman. A domain R is said to be t -linkative if each overring of R is t -linked over R . The class of t -linkative domains was introduced (but not named) in [11, Theorem 2.6] and named, in a detailed study, by the same authors and M. Roitman in [10]. They also introduced the notion of super- t -linkative domain, that is, a domain R such that each overring of R is t -linkative. Let $\mathcal{O}_t(R)$ denote the set of all t -linked overrings of R and $\mathcal{O}(R)$ the set of all overrings of R . Then $\mathcal{O}_t(R) \subseteq \mathcal{O}(R)$ and the inclusion may be strict (see [10, Example 4.1] for a domain R such that R' is not t -linked over R). Clearly an FO -domain is an FC -domain, an FC -domain is super- t -linkative ([23, Theorem 2.14] and [10, Corollary 2.5]) and a super- t -linkative domain is t -linkative. The t -linkative approach has proved its ability to be more efficient in the study of the set of overrings of integral domains. The purpose of this paper is to continue the investigation of some finiteness conditions on the set of t -linked overrings of an integral domain. In Section 2, we answer, in the positive, the following two questions:

- (i) If $\mathcal{O}_t(R)$ is finite, then is $\mathcal{O}(R)$ finite? and do we have $\mathcal{O}_t(R) = \mathcal{O}(R)$?
- (ii) If each chain of t -linked overrings of R is finite, then is R an FC -domain?

As FO -domains and FC -domains have finite spectrum, we list some results relating the (Krull) dimension of R to the cardinality of $\mathcal{O}(R)$. Section 3 is devoted to the study of Noetherian-like settings. The main result asserts that if a Mori domain R is an FC -domain, then its complete integral closure $\bar{R}$ is a Dedekind domain and the conductor $(R : \bar{R}) \neq (0)$, and so R is a one-dimensional domain. Moreover, if $A = (R : \bar{R})$ is a finitely generated ideal of R , then R is Noetherian (Theorem 3.6).

Let $SD(R)$ be the set of all (nonzero) strongly divisorial ideals of R (we recall that a nonzero ideal I is strongly divisorial if $I = II^{-1} = I_v$) and let $\phi_R : SD(R) \rightarrow \mathcal{O}_t(R)$, $I \mapsto (I : I) = I^{-1}$. Then ϕ_R is an injective map (note that ϕ_R is well-defined since $(I : I)$ is a t -linked overring of R , for each strongly divisorial ideal I of R [11, Proposition 2.2, (e)]). In Section 4, we give necessary and sufficient conditions for ϕ_R to be surjective and so bijective. This leads us to compute the number of strongly divisorial ideals for some classes of integral domains with finitely many overrings. As an application, we prove that if R is Mori and ϕ_R is surjective, then R is an FC -domain. We also characterize PVD (pseudo-valuation domain) R for which ϕ_R is surjective and compute its strongly divisorial ideals. The last section deals with domains R such that R is t -linked under all its overrings. We prove that a such domains are exactly the one-dimensional local domains. The section closes with the study of the transfer of this notion to the pullbacks in order to provide original examples.

Throughout, we denote by R' (resp. $\bar{R}$) the integral (resp. complete integral) closure of R and we use the symbol " $\subset$ " for the strict inclusion. For a nonzero (fractional) ideal I of R , $I^{-1} = (R : I) = \{x \in K | xI \subseteq R\}$. The v - and t -closures of I are defined, respectively, by $I_v = (I^{-1})^{-1}$ and $I_t = \bigcup J_v$, where J ranges over the set of finitely generated subideals of I . The ideal I is said to be a v -ideal (or

divisorial) if $I = I_v$, and a t -ideal if $I = I_t$. A t -maximal ideal is a t -ideal that is maximal for the inclusion. Finally, a local domain stands for a domain with exactly one maximal ideal and a semilocal domain is a domain with a finite number of maximal ideals. Unreferenced material is standard, typically as in [24].

2. GENERAL SETTINGS

We start this section by showing that if $O_t(R)$ is finite, then so is $O(R)$; and $O_t(R) = O(R)$. We also prove that if every chain of t -linked overrings is finite, then R is an FC -domain. This yields that the t -linked approach is more efficient than the Gilmer treatment of FO -domains and FC -domains (see [23]). As FO -domains and FC -domains have finite spectrum, so finite dimension, we list a few results treating the relation between the Krull dimension of an FO -domain R and the cardinality of $O(R)$.

Proposition 2.1. *If $O_t(R)$ is finite then $O(R) = O_t(R)$ and therefore R is an FO -domain.*

The proof need the following two results due to M. Zafrullah. For the convenience of the reader, we include them with their proofs [34].

Proposition 2.2. *Let $*$ be a star operation of finite type on R . If R has only a finite number of distinct maximal $*$ -ideals $P_1, \dots, P_r$, then $P_1, \dots, P_r$ are precisely the maximal ideals of R .*

Proof. Let $*$ be a star operation of finite type. Suppose that $P_1, \dots, P_r$ are the distinct maximal $*$ -ideals of R . Since for each nonzero nonunit $a \in R$ the ideal aR is a $*$ -ideal and so is must be contained in at least one maximal $*$ -ideal, we conclude that $R \setminus U \subseteq P_1 \cup P_2 \cup \dots \cup P_r$, where U is the group of units of R . Now, as each maximal ideal M consists of nonunits, we have $M \subseteq P_1 \cup P_2 \cup \dots \cup P_r$. By the prime avoidance lemma we have $M \subseteq P_i$ for some i . But since M is maximal, we have $M = P_i$. Next, as each P_j is contained in a maximal ideal which in turn is contained in some P_k and since P_i are incomparable, we conclude that each of P_i is a maximal ideal. $\square$

Corollary 2.3. *If R has a finite number of maximal t -ideals then every maximal ideal is a t -ideal.*

Proof of the Proposition 2.1 Assume that $O_t(R)$ is finite. Since for each t -maximal ideal M of R , R_M is t -linked over R , then R has only finitely many maximal t -ideals. By Corollary 2.3, every maximal ideal is a t -ideal. Hence $O(R) = O_t(R)$ by [11, Theorem 2.6]. $\square$

Proposition 2.4. *R is an FC -domain if and only if each chain of t -linked overrings of R is finite.*

Proof. Assume that each chain of t -linked overrings of R is finite. Then R has finitely many maximal t -ideals. Indeed, suppose on the contrary that R has infinitely many maximal t -ideals. Let $\{M_i\}_{i \geq 1}$ be an index ordered set of t -maximal ideals of R . For each $n \geq 1$ set $R_n = \bigcap_{i=1}^{i=n} R_{M_i}$. Then R_n is a t -linked overring of R (as an intersection of t -linked overrings, [11, Proposition 2.2, (b)]). Hence $\{R_n\}_{n \geq 1}$ is an infinite chain of t -linked overrings of R [23, Lemma 1.4], which is absurd. So

R has a finite number of t -maximal ideals. By Corollary 2.3 every maximal ideal is a t -ideal and by [11, Proposition 2.2], $O(R) = O_t(R)$. It follows that R is an FC -domain. $\square$

It's easy to see that for any domain R , $1 + \dim R \leq |O(R)| \leq |SSFc(R)|$, where $SSFc(R)$ is the set of all semistar operations of finite character on R . In particular, if $SSFc(R)$ is finite, then R is an FO -domain. The next theorem restates the following results [32, Theorem 7] and [31, Theorem 4.3 and Theorem 4.4] by substituting $O(R)$ to $SSFc(R)$. The proofs are the same with a minor changes. For the convenience of the reader, we include them here.

Theorem 2.5. *Let R be an integral domain of finite dimension.*

- (1) $|O(R)| = 1 + \dim R$ if and only if R is a valuation domain [32, Theorem 7].
- (2) $|O(R)| = 2 + \dim R$ if and only if R is a local domain, R' is a valuation domain and each proper overring of R contains R' . In this case $O(R) = \{R\} \cup O(R')$ [31, Theorem 4.4].
- (3) Assume that R is not local. Then $3 + \dim R \leq |O(R)|$, and the equality holds if and only if R is a Prüfer domain with exactly two maximal ideals and Y -graph spectrum [31, Theorem 4.3].

Proof. Set $n = \dim R$ and let $(0) \subset P_1 \subset \cdots \subset P_n$ be a chain of prime ideals of R such that $\dim R = ht P_n = n$ and let $\mathbb{E} = \{L, R_{P_1}, \dots, R_{P_n}\}$. Clearly $|\mathbb{E}| = n + 1$. Now, assume that $|O(R)| = 1 + \dim R$, then $O(R) = \mathbb{E}$, and therefore each overring of R is flat (as a localization of R). Hence R is a Prüfer domain. Clearly R is local with maximal ideal P_n (and so $R_{P_n} = R$). Otherwise, there is a maximal ideal M of R such that $M \neq P_n$. Then $O(R) = \mathbb{E} \cup \{R_M\}$, which is absurd. Hence R is local and therefore a valuation domain.

The converse is clear since the overrings of a valuation domain R are exactly the localizations of R at prime ideals.

(2) We claim that R is local. Indeed, set $n = \dim R$ and let $(0) \subset P_1 \subset \cdots \subset P_n = M$ be a chain of prime ideals of R such that $\dim R = ht M = n$. If N is a maximal ideal of R such that $N \neq M$, then $\{R, R_{P_1}, \dots, R_{P_{n-1}}, R_M, R_N, L\} \subseteq O(R)$. So $|O(R)| \geq 3 + n = 3 + \dim R$, which is absurd. Hence R is local. If R is integrally closed, then R is a valuation domain [23, Theorem 1.5] and so $|O(R)| = 1 + \dim R$ by Proposition 2.5, which is absurd. Hence $R \subset R'$ and therefore $1 + \dim R = 1 + \dim R' \leq |O(R')| \leq |O(R)| - 1 = 1 + \dim R = 1 + \dim R'$. So $1 + \dim R' = |O(R')|$ and therefore R' is a valuation domain. It follows that $O(R) = \{R, R_{P_1}, \dots, R_{P_{n-1}}, R', L\}$. Now, for each $i \in \{1, \dots, n-1\}$, let Q_i be a prime ideal of R' such that $Q_i \cap R = P_i$. Since $R'_{Q_i} \in O(R)$, then $R'_{Q_i} = R_{P_j}$ for some $j \in \{1, \dots, n-1\}$. Hence $Q_i R'_{Q_i} = P_j R_{P_j}$ and therefore $P_i = P_j$. So $i = j$ and therefore $R' \subseteq R'_{Q_i} = R_{P_i}$. Hence every proper overring of R contains R' and therefore $O(R) = O(R') \cup \{R\}$.

Conversely, assume that R' is a valuation domain and $O(R) = O(R') \cup \{R\}$. Then $|O(R)| = 1 + |O(R')| = 1 + (1 + \dim R') = 2 + \dim R' = 2 + \dim R$, as desired.

(3) Assume that R is not local. Set $n = \dim R$ and let $(0) \subset P_1 \subset \cdots \subset P_n = M$ be a chain of prime ideals of R such that $\dim R = ht M = n$. Let N be maximal ideal of R such that $N \neq M$ and let $\mathbb{E} = \{L, R_{P_1}, \dots, R_{P_{n-1}}, R_M, R_N, R\}$. Clearly $\mathbb{E} \subseteq O(R)$ and $|\mathbb{E}| = n + 3$. Hence $3 + \dim R \leq |O(R)|$. Now, assume that

$|O(R)| = 3 + \dim R = |\mathbb{E}|$. Then $O(R) = \mathbb{E}$. Since R is not a field and R' is integral over R , then R' cannot be a localization of R . Hence $R = R'$. Then R is an integrally closed FO -domain. By [23, Theorem 1.5], R is a Prüfer domain with finite spectrum, and clearly $\text{Spec}(R) = \{(0) \subset P_1 \subset \dots \subset P_n = M, N\}$. It remains to prove that $P_{n-1} \subseteq N$.

Suppose that N and P_{n-1} are not comparable. Set $T = R_N \cap R_{P_{n-1}}$. Then T is a quasilocal Prüfer domain with exactly two maximal ideals $Q_1 = NR_N \cap T$ and $Q_2 = P_{n-1}R_{P_{n-1}} \cap T$ (since T is the intersection of the two valuation domains R_N and $R_{P_{n-1}}$ that are not comparable, [8, Propositions 1& 2, page 412]). Clearly $Q_1 \neq M$ (otherwise, $N = M$, which is absurd) and $Q_2 \neq M$ (otherwise, $P_{n-1} = M$, which is absurd). Hence $R \neq T$. Since all the other overrings of R are valuation domains, then $O(R) = \mathbb{E} \subset \mathbb{E} \cup \{T\} \subseteq O(R)$, a contradiction. Hence N and P_{n-1} are comparable and by maximality $P_{n-1} \subset N$, as desired.

Conversely, assume that R is Prüfer with exactly two maximal ideals M and N and Y -graph spectrum, that is, $(0) \subset P_1 \subset \dots \subset P_{n-1} \subseteq M \cap N$ and let $\mathbb{E} = \{L, R_{P_1}, \dots, R_{P_{n-1}}, R_M, R_N, R\}$. Clearly $|\mathbb{E}| = 3 + n = 3 + \dim R$ and since each overring of R is a subintersection (i.e., intersection of localizations of R at some primes), then $O(R) = \mathbb{E}$, as desired. $\square$

For each positive integer $n \geq 1$, there exists an n -dimensional local domain R with $(3 + \dim R)$ overrings, as it's shown by the following example.

Example 2.6. Let k be a field and $X_1, \dots, X_{n+1}$ indeterminates over k . Set $R_1 = k[[X_1^2, X_1^5]]$. Clearly R_1 is a one-dimensional Noetherian local domain and it's easy to see that $O(R_1) = \{R_1, k[[X_1^2, X_1^3]], k[[X_1]] = R'_1, L_1 = qf(R_1) = k((X_1))\}$. So $|O(R_1)| = 4 = 3 + \dim R_1$.

Let $V_1 = L_1[[X_2]] = L_1 + M_1$, where $M_1 = X_2V_1$ and set $R_2 = R_1 + M_1$. Since R_1 is local, then so is R_2 and $\dim R_2 = \dim R_1 + \dim V_1 = 1 + 1 = 2$ [7, Theorem 2.1]. Since each overring of R_2 is comparable to V_1 [7, Theorem 3.1] and V_1 is a DVR , then $O(R_2) = \{T + M | T \in O(R_1)\} \cup \{L_2 = qf(R_2) = k((X_1, X_2))\}$. Therefore $|O(R_2)| = |O(R_1)| + 1 = 4 + 1 = 5 = 3 + \dim R_2$.

By induction on n , assume that R_n is a local domain with $\dim R_n = n$ and $|O(R_n)| = 3 + \dim R_n$. Let $L_n = qf(R_n)$ and set $V_n = L_n[[X_{n+1}]] = L_n + M_n$, where $M_n = X_{n+1}V_n$. Set $R_{n+1} = R_n + M_n$. Then $\dim R_{n+1} = \dim R_n + \dim V_n = n + 1$ and $|O(R_{n+1})| = |O(R_n)| + 1 = 3 + \dim R_n + 1 = 3 + \dim R_{n+1}$, as desired.

We end this section by the following result which shows that the sets $O_t(R)$ of all t -linked overrings of R and $O_w(R)$ of all w -overrings of a domain R are the same and the notions of t -linknative domain and DW -domain coincide. First we recall the following definitions.

- (i) An overring T of a domain R is said to be a w -overring of R if $T_w = T$ [13]. Let $\mathcal{O}_w(R)$ denote the set of all w -overrings of R .
- (ii) A domain R is said to be a DW -domain if each ideal of R is a w -ideal [30].

Proposition 2.7. *Let R be an integral domain. Then*

- (a) *Each w -overring of R is t -linked over R , that is, $\mathcal{O}_w(R) = \mathcal{O}_t(R)$.*
- (b) *The following assertions are equivalent*
 - (i) *R is t -linknative;*
 - (ii) *R is a DW -domain;*

Proof. The first part of this proposition was proved in [12, Proposition 3.1] and the second part appears in [33]. However, we give here a simple proof. Suppose that R is t -linkative. Let I be a nonzero ideal of R and let $x \in I_w$. Then there exists a f. g. ideal J of R such that $J^{-1} = R$ and $xJ \subseteq I$. By [11, Theorem 2.6], $J = R$. Hence $x \in I$ and therefore $I = I_w$, as desired. The converse follows also from [11, Theorem 2.6] since $I_t = R$ if and only if $I_w = R$. $\square$

3. NOETHERIAN-LIKE SETTINGS

Before starting this section, we recall the following useful definitions: An integral domain R is said to be:

- (1) Mori domain if R satisfies the *acc* condition on the v -ideals, and seminormal if $x \in R$, for each $x \in K$ with $x^2, x^3 \in R$ (see [5] for more details about Mori seminormal domains).
- (2) A conducive domain if $(R : T) \neq (0)$ for each overring T of R with $T \subsetneq K$ (see [9]).
- (3) Almost Krull if R_M is a Krull domain for each maximal ideal M of R , [24].
- (4) A pseudo-valuation domain (*PVD* for short) if there exists a valuation overring V of R such that $\text{Spec}(R) = \text{Spec}(V)$, [18, 19].
- (5) Quasi-Prüfer, if R' is Prüfer, [15, Corollary 6.5.14].

Proposition 3.1. *Let R be a Krull domain. The following statements are equivalent*

- (i) R is super- t -linkative;
- (ii) R is t -linkative;
- (iii) R is a Dedekind domain.

Proof. (i) $\implies$ (ii) Trivial.

(ii) $\implies$ (iii) Assume that R is t -linkative. By [11, Theorem 2.6] each t -invertible ideal is invertible. Since R is Krull, then each nonzero ideal is t -invertible, and so invertible. Hence R is Dedekind.

(iii) $\implies$ (i) Follows from [10, Proposition 3.13]. $\square$

Corollary 3.2. *Let R be an almost Krull domain. Then R is super- t -linkative if and only if R is almost Dedekind.*

Proof. Let M be a maximal ideal of R . Then R_M is a Krull domain. Since R_M is super- t -linkative, then R_M is a Dedekind domain. Hence R is almost Dedekind. Conversely, if R is almost Dedekind, then R is a Prüfer domain. By [10, Corollary 2.5], R is super- t -linkative. $\square$

In [10, Proposition 3.13], it was proved that a Noetherian domain R is super- t -linkative if and only if $\dim R \leq 1$. Our next two results show that a one-dimensional Mori domain needs not be super- t -linkative, and state conditions under which a super- t -linkative Mori domain is of dimension one.

Lemma 3.3. *Let R be a Mori domain. If R is t -linkative, then every prime ideal is divisorial.*

Proof. Let P be a prime ideal of R . If $htP = 1$, then P is divisorial, [5, Theorem 3.1], as desired. Assume that $htP \geq 2$. Also by [5, Theorem 3.1], either P is

strongly divisorial or $P^{-1} = R$. If $P^{-1} = R$, since R is Mori, then $P_t = P_v = R$, which contradicts [11, Theorem 2.6]. Hence P is divisorial, as desired. $\square$

Proposition 3.4. *Let R be a Mori domain such that either (i) $(R : \bar{R}) \neq (0)$, or (ii) R is seminormal. If R is super- t -linkative, then $\dim R = 1$.*

Proof. (i) Assume that $(R : \bar{R}) \neq (0)$. Then $\bar{R}$ is a Krull domain, [4, Corollary 18] or [5, Theorem 7.4]. Since $\bar{R}$ is super- t -linkative (as an overring of R), by Proposition 3.1, $\bar{R}$ is a Dedekind domain. So $\dim \bar{R} = 1$. By [6, Corollary 3.4], $\dim R = 1$.

(ii) Assume that R is seminormal. Suppose that $\dim R \geq 2$ and let M be a maximal ideal of R such that $ht M \geq 2$. Since R_M is a super- t -linkative Mori domain which is seminormal, then without loss of generality, we may assume that R is local with maximal ideal M . Since $ht M \geq 2$, by [6, Lemma 2.5], $B = (M : M)$ contains a nondivisorial prime ideal Q such that $Q \cap R = M$. Since $ht M \geq 2$, then $M^{-1} = (M : M)$. So B is a Mori domain which is super- t -linkative and which contains a non divisorial prime ideal, which is absurd by Lemma 3.3. It follows that $\dim R = 1$. $\square$

The converse is not true as it's shown by the following example. First we recall that the valuative dimension of R , denoted by $\dim_v R$, is given by $\dim_v R = \max\{\dim V \mid V \text{ valuation overring of } R\}$.

Example 3.5.

Let k be a field and X, Y and Z indeterminates over k . Set $R = k + Zk(X, Y)[[Z]] = k + M$, where $M = Zk(X, Y)[[Z]]$. Then R is an integrally closed Mori domain (in fact R is a PVD with maximal ideal M and associated valuation overring $V = k(X, Y)[[Z]]$). $\bar{R} = V = k(X, Y)[[Z]]$, $(R : \bar{R}) = M$. Since $\dim_v R = 3$, by [10, Proposition 3.12], R is not super- t -linkative.

From [23, Theorem 2.14], it's easy to see that a Noetherian domain R is an FC -domain if and only if $R' = \bar{R}$ is a quasilocal Dedekind domain and $(R : R') \neq (0)$.

Theorem 3.6. *Let R be a Mori domain. If R is an FC -domain, then $\bar{R}$ is a semilocal Dedekind domain and $(R : \bar{R}) \neq (0)$, in particular $\dim_v R = 1$. Moreover, if $A = (R : \bar{R})$ is a finitely generated ideal of R , then R is Noetherian.*

Proof. Since every FC -domain is super- t -linkative, by Proposition 3.4, it suffices to show that $(R : \bar{R}) \neq (0)$. By [23, Theorem 2.14], R' is a Prüfer domain and R' is a finite R -module. So the conductor $A = (R : R') \neq (0)$. Set $T = (A_v : A_v) = (AA^{-1})^{-1}$. Then T is a Mori domain. Since A is an ideal of R' , then $R' \subseteq (A : A) \subseteq (A_v : A_v) = T$. Since R' is Prüfer, then so is T . Hence T is a Dedekind domain (as a Prüfer domain which is also Mori). Clearly $(R : T) = (AA^{-1})_v \neq (0)$ and so R and T have the same complete integral closure, that is, $\bar{R} = \bar{T} = T$. Hence $\bar{R}$ is Dedekind and so $\dim \bar{R} = 1$. Since $\bar{R}$ is an FC -domain (as an overring of R), then $\bar{R}$ has finite spectrum. So $\bar{R}$ is semilocal, and clearly $(R : \bar{R}) = (R : T) = (AA^{-1})_v \neq (0)$, as desired. Since $\dim R = \dim \bar{R} = 1$, [6, Corollary 3.4. (1)], and R' is a Prüfer domain, then $\dim_v R = \dim_v R' = \dim R' = \dim R = 1$. Now, assume that A is a finitely generated ideal of R . Let P be a nonzero prime ideal of R . Since

$\dim R = 1$, then $P = M$ is a maximal ideal which is divisorial. If $MM^{-1} = R$, then M is finitely generated (as an invertible ideal). Assume that $MM^{-1} \subset R$. Then $MM^{-1} = M$. Hence $M^{-1} = (M : M) \subseteq \bar{R} = A^{-1}$. So $A = A_v \subseteq M_v = M$. By [23, Theorem 2.14], R/A is Artinian. Hence M/A is a finitely generated ideal of R/A . Set $M/A = (\bar{x}_1, \dots, \bar{x}_n)$, where $x_i \in M$ for each $i = 1, \dots, n$. Let $J = (x_1, \dots, x_n)$. Clearly $M = J + A$, and so M is finitely generated. Hence every prime ideal of R is finitely generated and therefore R is Noetherian. $\square$

We recall that a domain R is said to be quasi-coherent Mori domain if every t -ideal is finitely generated [29, page 85].

Corollary 3.7. *Let R be a quasi-coherent Mori domain. If R is an FC-domain, then R is Noetherian.*

Proof. Since $A = (R : \bar{R})$ is a v -ideal of R , then A is finitely generated. The conclusion follows from Theorem 3.6. $\square$

According to [13], a domain R is said to be strong Mori (SM for short) if R satisfies the ascending chain conditions on w -ideals. Noetherian domains are strong Mori and strong Mori domains are Mori.

Corollary 3.8. *Let R be a strong Mori domain. If R is an FC-domain, then R is a one-dimensional Noetherian domain.*

Proof. Follows immediately from Theorem 3.6 and [13, Corollary 1.10]. $\square$

4. t -LINKED OVERRINGS AND STRONGLY DIVISORIAL IDEALS

We recall that a nonzero ideal I of a domain R is said to be strong (or a trace ideal) if $I = II^{-1}$ and strongly divisorial if it is strong and divisorial, that is $I = I_v = II^{-1}$. Let $SD(R)$ denotes the set of all nonzero strongly divisorial ideals of R . By [11, Proposition 2.2], for each ideal A of a domain R , $(A_v : A_v)$ is t -linked over R . In particular, if $I \in SD(R)$, then $I^{-1} = (I : I)$ is t -linked over R . This yields an injection map $\phi_R : SD(R) \longrightarrow \mathcal{O}_t(R) \setminus \{L\}$, $I \mapsto I^{-1}$ (we note that this map was introduced by V. Barucci in [4]). The following proposition characterizes when ϕ_R is surjective.

Proposition 4.1. *Let R be a domain.*

- 1) ϕ_R is surjective if and only if each t -linked overring of R is a fractional v -ideal of R .
- 2) If ϕ_R is surjective, then:
 - i) R is a conducive domain;
 - ii) Every non maximal prime ideal is strong.
 - iii) If R is not local, then every maximal ideal of R is not divisorial, that is, for each maximal ideal M of R , $M^{-1} = R$.

Proof. 1) $\implies$ Let $T \in \mathcal{O}_t(R) \setminus \{L\}$. Then there exists $I \in SD(R)$ such that $T = I^{-1}$. Hence T is a fractional v -ideal of R .

$\impliedby$ Let $T \in \mathcal{O}_t(R)$ and set $I = (R : T)$. Since T is a fractional v -ideal of R , then $T \subseteq (I : I) \subseteq I^{-1} = T_v = T$. Hence $I \in SD(R)$ and $\phi_R(I) = T$, as desired.

2) i) Since R is not a field, then $\dim R \geq 1$. Let P be a t -prime ideal of R . Now, for each overring $T \subset L$ of R , $T_{R \setminus P}$ is t -linked over R [11, Proposition 2.9].

Since ϕ_R is surjective, then there is $I \in SD(R)$ such that $I^{-1} = T_{R \setminus P}$. Hence $I = I_v = (R : I^{-1}) = (R : T_{R \setminus P}) \subseteq (R : T)$. So $(R : T) \neq (0)$ and therefore R is conducive.

ii) Let P be a nonmaximal prime ideal of R . Since $R_P \in O_t(R) \setminus \{L\}$, then $R_P = I^{-1}$ for some $I \in SD(R)$. Hence $IR_P = II^{-1} = I$. Since P is not maximal, then $R \subset R_P = I^{-1}$. Hence $I \subseteq P$. So $P^{-1} \subseteq I^{-1} = R_P$ and then $PP^{-1} \subseteq PR_P$. Since $PP^{-1} \subseteq R$, then $PP^{-1} \subseteq PR_P \cap R = P$. Hence $PP^{-1} = P$, as desired.

iii) Assume that R is not local. Let M be a maximal ideal of R and $I \in SD(R)$ such that $I^{-1} = R_M$. Then $I = II^{-1} = IR_M$. Since $R \subset R_M$, then $I \subseteq M$. So $M^{-1} \subseteq I^{-1} = R_M$. Since $M^{-1} \subseteq R_N$ for each maximal ideal $N \neq M$, then $M^{-1} \subseteq R_M \cap (\cap_{N \neq M} R_N) = R$. Hence $M^{-1} = R$, as desired. $\square$

Corollary 4.2. *Let R be a Mori domain. If ϕ_R is surjective, then R is a one-dimensional conducive local domain and either (i) R is a DVR, or (ii) the maximal ideal M of R is strongly divisorial and each proper overring of R contains M^{-1} .*

Proof. By Proposition 4.1, R is conducive. So $\bar{R}$ is a one-dimensional valuation domain (it easy to see that the only overrings of $\bar{R}$ are $\bar{R}$ and $L = qf(R)$). Since $(R : \bar{R}) \neq (0)$, then $\dim R = 1$ [6, Corollary 3.4]. Since R is Mori, then $\bar{R}$ is Krull and therefore $\bar{R}$ is a DVR. Set $\text{Spec}(\bar{R}) = \{(0) \subset N\}$. Now, if Q and M are maximal ideals of R , since $htQ = htM = \dim R = 1$, by [6, Proposition 1.1], there is Q' and M' in $\text{Spec}(\bar{R})$ such that $Q' \cap R = Q$ and $M' \cap R = M$. Since $\text{Spec}(\bar{R}) = \{(0) \subseteq N\}$, then $Q' = M' = N$ and therefore $Q = M$. Hence R is local with maximal ideal $M = N \cap R$.

1) If $MM^{-1} = R$, then R is a DVR.

2) If $MM^{-1} = M$, then set $T = M^{-1} = (M : M)$. Clearly T is a proper overring of R (since M is a v -ideal of R). Let S be a proper overring of R . Since $\dim R = 1$, then $O(R) = O_t(R)$. So S is t -linked over R . But, since ϕ_R is surjective, then there is $I \in SD(R)$ such that $S = I^{-1}$. Since S is a proper overring of R , then $I \subset R$. So $I \subseteq M$ (since R is local with maximal ideal M). Hence $T = M^{-1} \subseteq I^{-1} = S$, as desired. $\square$

Corollary 4.3. *Let R be a Mori domain. If ϕ_R is surjective, then R is an FC-domain.*

Proof. By Corollary 4.2, R is conducive and $\dim R = 1$. So $O(R) = O_t(R)$. Set $A = (R : \bar{R}) \neq (0)$ and let $\{R_n\}_{n \geq 1}$ be a chain of overrings of R . Since ϕ_R is surjective, then for each $n \geq 1$, there exists $I_n \in SD(R)$ such that $I_n^{-1} = R_n$. So for each $n \geq 1$, $R_n = I_n^{-1} = (I_n : I_n) \subseteq \bar{R} = A^{-1}$. Hence $A = A_v \subseteq (I_n)_v = I_n$. So $A \subseteq \bigcap_{n \geq 1} I_n$ and therefore $\bigcap_{n \geq 1} I_n \neq (0)$. Since $I_n^{-1} = R_n \subseteq R_{n+1} = I_{n+1}^{-1}$, then $I_{n+1} = (I_{n+1})_v \subseteq (I_n)_v = I_n$. Hence $\{I_n\}_{n \geq 1}$ is a decreasing chain of v -ideals of R with nonzero intersection. Since R is Mori, then $\{I_n\}_{n \geq 1}$ stabilizes, and therefore $\{R_n\}_{n \geq 1}$ stabilizes. It follows that R is an FC-domain. $\square$

Proposition 4.4. 1) *Let R be a valuation domain. Then ϕ_R is surjective. Moreover, if $\dim R = n \geq 1$ is finite, then R has exactly n strongly divisorial ideals.*
 2) *If R is completely integrally closed, then ϕ_R is surjective if and only if R is a one-dimensional valuation domain.*

3) If R is a Prüfer domain, then ϕ_R is surjective if and only if R is a valuation domain.

Proof. 1) Assume that R is a valuation domain. Then $O_t(R) = O(R)$. Let $T \in O(R) \setminus \{L\}$. If $T = R$, then $T = \phi_R(R)$. Assume that $R \subset T$. Then $T = R_Q$ for some nonzero nonmaximal prime ideal Q of R . By [22, Theorem 3.8 and Proposition 3.10], $Q \in SD(R)$ and $\phi_R(Q) = R_Q = T$. Moreover, if $\dim R = n$ is finite, then $SD(R) = \{\text{all nonzero nonmaximal prime ideals of } R\} \cup \{R\}$.

2) Assume that R is completely integrally closed and ϕ_R is surjective. It's clear that $SD(R) = \{R\}$. Since ϕ_R is surjective, then $O(R) = \{R, L\}$ and therefore R is one-dimensional valuation domain.

3) Assume that R is Prüfer and ϕ_R is surjective. Clearly $O(R) = O_t(R)$ since the t -operation on a Prüfer domain is trivial. It suffices to prove that R is local. If not, let M be a maximal ideal of R . Since ϕ_R is surjective, then there exists a proper ideal $I \in SD(R)$ such that $I^{-1} = R_M$. Since $IR_M = II^{-1} = I$, then $I \subseteq M$. Let N be a maximal ideal of R such that $N \neq M$. If $I \not\subseteq N$, then $R_M = I^{-1} \subseteq R_N$. So $N \subseteq M$ and hence $M = N$, which is absurd. It follows that $I \subseteq N$. Therefore I is contained in all maximal ideals of R . Now, let P be a minimal prime over I . By [22, Theorem 3.2], $R_M = I^{-1} \subseteq R_P$. Hence $P \subseteq M$. Since R is a Prüfer domain, then $\text{Spec}(R)$ is treed. So $\text{Min}(I)$, the set of all minimal primes over I , is reduced to one element, say $\text{Min}(I) = \{P\}$. Let N be a maximal ideal of R . Since $I \subseteq N$ and $\text{Min}(I) = \{P\}$, then $P \subseteq N$. We claim that $I^{-1} = R_P$. Indeed, by [22, Theorem 3.2], $I^{-1} \subseteq R_P$. Conversely, if there exists $x \in R_P$ and $a \in I$ such that $xa \notin R$, then $xa \notin R_N$ for some maximal ideal N of R . Since R_N is a valuation domain, then $(xa)^{-1} \in R_N$. Since $P \subseteq N$, then $R_N \subseteq R_P$. So $(xa)^{-1} \in R_P$. Hence $1 = (xa)^{-1}(xa) \in IR_P \subseteq PR_P$, which is absurd. It follows that $R_M = I^{-1} = R_P$ and therefore $P = M$. Hence M is the unique minimal prime over I and M is contained in all maximal ideals of R , which is a contradiction. It follows that R is local and therefore R is a valuation domain. $\square$

Theorem 4.5. *Let R be a PVD domain, V its associated valuation overring, M its maximal ideal, $k = R/M$ its residue field, $K = V/M$, and suppose that $R \subset V$. The following statements are equivalent.*

- i) ϕ_R is surjective;
- ii) The extension $R \subset V$ is minimal, i.e. there are no rings properly between R and V ;
- iii) K is algebraic over k and for each $\alpha \in K \setminus k$, $K = k(\alpha)$;
- iv) $O(R) = \{R\} \cup O(V)$.

Moreover, if $\dim V = n$ is finite, then R is an FO-domain and R has exactly $n+1$ strongly divisorial ideals $\{P_1, P_2, \dots, P_{n-1}, M, R\}$, where $P_1, P_2, \dots, P_{n-1}, M$ are the nonzero prime ideals of R .

Proof. We first note that $O_t(R) = O(R)$ since R is local with maximal ideal M and $M^{-1} = V$.

i) $\implies$ ii) Assume that ϕ_R is surjective. Let T be an overring of R such that $R \subset T \subseteq V$. By i) there exists $I \in SD(R)$ such that $I^{-1} = T$. Since $R \subset T$, then $I \subset R$ and since R is local, then $I \subseteq M$. So $V = M^{-1} \subseteq I^{-1} = T \subseteq V$. Hence

$V = T$ and therefore the extension $R \subset V$ is minimal.

$ii) \implies iii)$ Let $\alpha \in K \setminus k$ and set $T = \psi^{-1}(k[\alpha])$. Since $R \subset T \subseteq V$, by $ii)$, $T = V$. So $k[\alpha] = K$, as desired.

$iii) \implies iv)$ It's well-known that each overring of R is comparable to V . So $O(R) = [R, V[\cup O(V)$, where $[R, V[$ is the set of all overrings T of R such that $R \subseteq T \subset V$. Hence $O(R) = \{R\} \cup O(V)$ since $[R, V[= \{R\}$.

$iv) \implies i)$ Assume that $O(R) = \{R\} \cup O(V)$. Clearly $R = \phi_R(R)$, $V = M^{-1} = \phi_R(M)$ and $M \in SD(R)$. Let T be an overring of R . By $iv)$ and without loss of generality, we may assume that $T = V_P$, where P is a nonmaximal prime ideal of V . Since R and V have the same spectrum, P is a prime ideal of R and $V_P = (P : P) = P^{-1}$. So $P_v \in SD(R)$ and $T = V_P = (P_v : P_v) = \phi_R(P_v)$. It follows that ϕ_R is surjective. $\square$

It's clear that any $(n-1)$ -dimensional valuation domain, $n \geq 2$, has exactly n overrings. Also it's easy to see that a domain R has exactly two overrings if and only if R is one-dimensional valuation domain. The above theorem leads us to construct a PVD domain R_n which is not a valuation domain, and which has exactly n overrings, for any positive integer $n \geq 3$.

Example 4.6. Let n be a positive integer with $n \geq 3$, $\mathbb{Q}$ be the field of rational numbers and $X_1, \dots, X_{n-2}$ indeterminates over $\mathbb{Q}$. Our aim is to construct a descending chain of valuation domains $V_1 \supsetneq V_2 \supsetneq \dots \supsetneq V_{n-2}$ with maximal ideals $M_1 \subset M_2 \subset \dots \subset M_{n-2}$ (respectively) such that $\dim V_i = i$ for each $i \in \{1, \dots, n-2\}$ and $V_{n-2} = \mathbb{Q}(\sqrt{2}) + M_{n-2}$.

- For $n = 3$, just set $V_1 = \mathbb{Q}(\sqrt{2})[[X_1]] = \mathbb{Q}(\sqrt{2}) + M_1$, where $M_1 = X_1 V_1$.
- For $n = 4$, set $V_1 = \mathbb{Q}(\sqrt{2})((X_1))[[X_2]] = \mathbb{Q}(\sqrt{2})((X_1)) + M_1$, where $M_1 = X_2 V_1$, and $V_2 = \mathbb{Q}(\sqrt{2})[[X_1]] + M_1 = \mathbb{Q}(\sqrt{2}) + M_2$, where $M_2 = X_1 V_2$.
- For $n = 5$, set $V_1 = \mathbb{Q}(\sqrt{2})((X_1, X_2))[[X_3]] = \mathbb{Q}(\sqrt{2})((X_1, X_2)) + M_1$, where $M_1 = X_3 V_1$, $V_2 = \mathbb{Q}(\sqrt{2})((X_1))[[X_2]] + M_1 = \mathbb{Q}(\sqrt{2})((X_1)) + M_2$, where $M_2 = X_2 V_2$, and $V_3 = \mathbb{Q}(\sqrt{2})[[X_1]] + M_2 = \mathbb{Q}(\sqrt{2}) + M_3$, where $M_3 = X_1 V_3$.

Iterating this process, we construct the desired chain as follows:

Let $V_1 = \mathbb{Q}(\sqrt{2})((X_1, \dots, X_{n-3}))[[X_{n-2}]] = \mathbb{Q}(\sqrt{2})((X_1, \dots, X_{n-3})) + M_1$, where $M_1 = X_{n-2} V_1$.

$V_2 = \mathbb{Q}(\sqrt{2})((X_1, \dots, X_{n-4}))[[X_{n-3}]] + M_1 = \mathbb{Q}(\sqrt{2})((X_1, \dots, X_{n-4})) + M_2$, where $M_2 = X_{n-3} V_2$.

$V_3 = \mathbb{Q}(\sqrt{2})((X_1, \dots, X_{n-5}))[[X_{n-4}]] + M_2 = \mathbb{Q}(\sqrt{2})((X_1, \dots, X_{n-5})) + M_3$, where $M_3 = X_{n-4} V_3$.

$V_i = \mathbb{Q}(\sqrt{2})((X_1, \dots, X_{n-i-2})) + M_i$, where $M_i = X_{n-i-1} V_i$. Now, let $R_n = \mathbb{Q} + M_{n-2}$. Then:

$|O(R_n)| = 1 + |O(V_{n-2})| = 1 + (1 + \dim V_{n-2}) = 1 + 1 + n - 2 = n$, as desired.

5. DOMAINS THAT ARE t -LINKED UNDER ITS OVERRINGS

According to [2], A domain R is said to be t -linked under T if for each finitely generated ideal I of R , $(T : IT) = T$ implies that $I^{-1} = R$. In some sense the

notion of “ t -linked under” is the opposite of the t -linkdness. It was introduced by Anderson and Zafrullah in [2] to characterize almost Bézout domains. In this short section, we will prove some results related to this notion. First we note that if $T = L$, then $(L : IL) = L$ for every nonzero ideal I of R , however I^{-1} needs not be equal to R . It turns that if R is t -linked under its quotient field L , then $R = L$. In this view, and in accordance with our hypothesis in the introduction we always assume that $R \subseteq T \subset L$.

Proposition 5.1. *Let R be an integral domain and Q a nonzero prime ideal of R . If R is t -linked under R_Q , then R is local with maximal ideal Q .*

Proof. Let Q be a nonzero prime ideal of R and let $a \in R \setminus Q$. Set $I = aR$. Since $(R_Q : IR_Q) = R_Q$ and R is t -linked under R_Q , then $a^{-1}R = I^{-1} = R$. Then $a^{-1} \in R$ and therefore R is local with maximal ideal Q . $\square$

Proposition 5.2. *Let R be an integral domain. Then R is t -linked under each overring T of R if and only if R is a one-dimensional local domain.*

Proof. By Proposition 5.1, R is local with maximal ideal M , $\dim R = 1$ and $\text{Spec}(R) = \{(0) \subset M\}$.

Conversely, assume that R is local with maximal ideal M and $\dim R = 1$. If $|O(R)| = 2$, then R is the only overring of R properly contained in L , and trivially R is t -linked under itself. Assume that $|O(R)| > 2$. Let T be an overring of R which is not a field and let Q a prime t -ideal of T (such a prime t -ideal exists since T is not a field and it suffices to consider any minimal prime over a proper principal ideal). Since R is local and $\dim R = 1$, then $Q \cap R = M$. Now, let I be a finitely generated ideal of R such that $(T : IT) = T$. If $R \subset I^{-1}$, then I_v is a proper ideal of R . So $I_v \subseteq M$. Hence $IT \subseteq I_v T \subseteq MT \subseteq Q$. Therefore $T = (IT)_{t_1} \subseteq Q_{t_1} = Q$ (here t_1 denotes the t -operation with respect to T), which is absurd since Q is t -ideal. Hence $I^{-1} = R$ and therefore R is t -linked under T . $\square$

Proposition 5.3. *Let R be a Prüfer domain. If T is an overring of R such that R is t -linked under T , then $R = T$.*

Proof. Let T be an overring of R such that $T \subset L$. Assume that R is t -linked under T . By [24, Theorem 26.1] or [15, Theorem 1.1.2], $T = \bigcap \{R_P \mid P \in \Omega\}$, where Ω is the set of all prime ideals P of R such that $PT \subset T$. Let Q be a nonzero prime ideal of R .

If $QT = T$, then $1 = \sum_{i=1}^{i=n} a_i x_i$ where $a_i \in Q$ and $x_i \in T$ for each $i \in \{1, \dots, n\}$. Set

$I = \sum_{i=1}^{i=n} a_i R$. Then I is a finitely generated ideal of R and $IT = T$. So $(T : IT) = T$.

Since R is t -linked under T , then $I^{-1} = (R : I) = R$. But since R is a Prüfer domain, then I is invertible. So $R = II^{-1} = IR = I$ and hence $R = I \subseteq Q$, which is a contradiction. Hence $QT \subset T$. So $Q \in \Omega$ and therefore $\Omega = \text{Spec}(R) \setminus \{(0)\}$. It follows that $T = \bigcap \{R_P \mid P \in \Omega\} = \bigcap \{R_P \mid P \in \text{Spec}(R) \setminus \{(0)\}\} = R$, as desired. $\square$

Proposition 5.4. *Let R be an integral domain. If R' is Prüfer or $\dim R = 1$, then R is t -linked under R' .*

Proof. If $R = R'$, nothing to prove. So we may assume that $R \subset R'$. Let I be a nonzero finitely generated ideal of R such that $(R' : IR') = R'$ and suppose that

$I \subset R$. Let P be a prime ideal of R such that $I \subseteq P$ and let Q be a prime ideal of R' such that $Q \cap R = P$ (such a prime ideal exists since R' is integral over R , so the extension $R \subset R'$ is Lying Over).

Assume that R' is Prüfer. Since IR' is a finitely generated ideal, then $R' = IR'(R' : IR') = IR' \subseteq PR' \subseteq Q$, which is absurd. Hence $I = R$ and therefore $I^{-1} = R$, as desired.

Now, if $\dim R = 1$. Then $\dim R' = 1$. So $ht Q = 1$ and therefore Q is a t -prime ideal of R' . But $(R' : IR') = R'$ implies that $R' = (IR')_{v'} = (IR')_t \subseteq (Q)_{t'} = Q$, where t' and v' are the t - and v -operations with respect to R' . This yields a contradiction. Hence, also in this case, $I = R$ and therefore $(R : I) = R$, as desired. $\square$

It's not the case that R is always t -linked under R' as it's shown by the following example.

Example 5.5. Let $\mathbb{Q}$ be the field of rational numbers and X and Y indeterminates over $\mathbb{Q}$. Set $T = \mathbb{Q}(\sqrt{2})[[X, Y]] = \mathbb{Q}(\sqrt{2}) + M$ and let $R = \mathbb{Q} + M$. Clearly R is a Noetherian domain which is local with maximal ideal M and $\dim R = 2$ [17, Theorem 4.12]. Since $(R : T) = M$, then $R' = \bar{R} = \bar{T} = T$. However, R is not t -linked under T since M is a finitely generated ideal of R , $(T : MT) = (T : M) = T$ and $M^{-1} = T$ (or by Theorem 5.6 below, since M is not a t -ideal of T).

In the literature, a few examples satisfying this notion are known. As pullbacks are known as a source for examples and counterexamples, the next results characterize this notion under pullbacks.

Let T be an integral domain which is not a field, M a maximal ideal of T , $K = T/M$ its residue field, $\phi : T \rightarrow K = T/M$ the canonical projection and D a subring of K . Let R be the pullback of the following diagram

$$\begin{array}{ccc} R & \longrightarrow & D \\ \downarrow & & \downarrow \\ T & \xrightarrow{\phi} & K = T/M \end{array}$$

We assume that $R \subset T$ and we refer to this diagram as diagram of type $(\square)$.

Theorem 5.6. *For the diagram of type $(\square)$, R is t -linked under T if and only if $D = k$ is a field and M is a t -ideal of T .*

Proof. Let $0 \neq d \in D$, $J = dD$ and $I = \phi^{-1}(J)$. By [14, Corollary 1.7], I is a finitely generated ideal of R . Since $M \subset I$ and M is a maximal ideal of T , then $IT = T$. Hence $(T : IT) = T$. Since R is t -linked under T , then $I^{-1} = R$. By [21, Proposition 6], $d^{-1}D = J^{-1} = D$. Hence $d^{-1} \in D$ and therefore D is a field.

Let t_1 denote the t -operation on T . Suppose that M is not a t -ideal of T .

Then $M_{t_1} = T$. So there exists a finitely generated subideal $A = \sum_{i=1}^{i=r} a_i T$ of M

such that $(T : A) = T$. Set $B = \sum_{i=1}^{i=r} a_i R$. Then B is a f.g. subideal of M with

$(T : BT) = (T : A) = T$. Since R is t -linked under T , then $B^{-1} = R$. Hence $R = B_v \subseteq M_v = M$ [21, Corollary 2], which is absurd. It follows that M is a t -ideal of T .

Conversely, assume that $D = k$ is a field and M is a t -maximal ideal of T . Let I be a f.g. ideal of R such that $(T : IT) = T$. Then $I^{-1} = (R : I) \subseteq (T : IT) = T$ and so $M = (R : T) \subseteq I_v = I_t$. Since $D = k$ is a field, then M is a maximal ideal of R . Hence either $M = I_v = I_t$ or $I_v = I_t = R$. Suppose that $M = I_v$. Then $I \subseteq M$. So $IT \subseteq M$ and therefore $T = (IT)_{v_1} = (IT)_{t_1} \subseteq M_{t_1} = M$, which is absurd. Hence $I_v = R$ and therefore $I^{-1} = R$. It follows that R is t -linked under R . $\square$

Corollary 5.7. (1) *Let R be a PVD (pseudo-valuation domain) and V its associated valuation overring. Then R is t -linked under V .*

(2) *Let $R = k + XK[X]$, where $k \subset K$ is an extension of fields. Then R is t -linked under $K[X]$.*

REFERENCES

- [1] D. D. Anderson, D. Dobbs and B. Mullins, *The primitive element theorem for commutative algebras*, Houston J. Math. **25** (1999) 603–623.
- [2] D. D. Anderson and M. Zafrullah, *Almost Bézout domains*, J. Algebra **142** (1991) 285–309.
- [3] A. Ayache and A. Jaballah, *Residually algebraic pairs of rings*, Math. Zeit. **225** (1997) 49–65.
- [4] V. Barucci, *Strongly divisorial ideals and complete integral closure of an integral domain*, J. Algebra **99** (1986) 132–142.
- [5] V. Barucci, *Mori domains*, in “Non-Noetherian Commutative Ring Theory” Mathematics and its Applications, Kluwer (S. Chapman and S. Galz ed.) Vol.520 (2000) 57–73.
- [6] V. Barucci and E. Houston, *On the prime spectrum of a Mori domain*, Comm. Algebra **24** (1996) 3599–3622.
- [7] E. Bastida and R. Gilmer, *Overrings and divisorial ideals of rings of the form $D+M$* , Michigan Math. J. **20** (1973) 79–95.
- [8] N. Bourbaki, *Elements of Mathematics, Commutative Algebra*, Hermann, 1972.
- [9] D. Dobbs and R. Fedder, *Conducive integral domains*, J. Algebra **86** (1984) 494–510.
- [10] D. Dobbs, E. Houston, T. Lucas, M. Roitman and M. Zafrullah, *On t -linked overrings*, Comm. Algebra **20** (1992) 1463–1488.
- [11] D. Dobbs, E. Houston, T. Lucas and M. Zafrullah, *t -linked overrings and Prüfer v -multiplication domains*, Comm. Algebra **17** (1989) 2835–2852.
- [12] G. W. Chang, *Strong Mori domains and the ring $D[X]_{N_v}$* , J. Pure Appl. Algebra **197** (2005) 279–292.
- [13] W. Fagui and R. L. McCasland, *On strong Mori domains*, J. Pure Appl. Algebra **135** (1999) 155–165.
- [14] M. Fontana and S. Gabelli, *On the Class group and the Local Class Group of a Pullback*, J. Algebra **181** (1996) 803–835.
- [15] M. Fontana, J. Huckaba and I. Papick, *Prüfer domains* Marcel Dekker, **203** New York, 1997.
- [16] M. Fontana, J. Huckaba, I. Papick and M. Roitman, *Prüfer domains and endomorphisms of their ideals*, J. Algebra **157** (1993) 489–516.
- [17] S. Gabelli and E. Houston, *Coherentlike Conditions in Pullbacks*, Michigan Math. J. **44** (1997) 99–122.
- [18] J. Hedstrom and E. Houston, *Pseudo-valuation domains*, Pacific J. Math. **75** (1978) 137–147.
- [19] J. Hedstrom and E. Houston, *Pseudo-valuation domains II*, Houston J. Math. **4** (1978) 199–207.
- [20] W. Heinzer and I. Papick, *The Radical Trace Property*, J. Algebra **112** (1988) 110–121.
- [21] E. Houston, S. Kabbaj, T. Lucas and A. Mimouni, *Duals of Ideals in Pullback Constructions*, Lecture Notes Pure Appl. Math. Dekker **171** (1995) 263–276.
- [22] J. A. Huckaba and I. J. Papick, *When the dual of an ideal is a ring*, Manuscripta Math. **37** (1982) 67–85.
- [23] R. Gilmer, *Some finiteness conditions on the set of overrings of an integral domain*, Proc. Amer. Math. Soc. **131** (2002) 2337–2346.
- [24] R. Gilmer, *Multiplicative ideal theory*, Pure and Applied Mathematics, No. 12. Marcel Dekker, Inc., New York, 1972.

- [25] A. Jaballah, *A lower bound for the number of intermediate rings*, Comm. Algebra **27** (1999) 1307–1311.
- [26] A. Jaballah, *Finiteness of the set of intermediary rings in normal pairs*, Saitama Math. J. **17** (1999) 59–61.
- [27] A. Jaballah, *The number of overrings of an integrally closed domain*, Expo. Math. **23** (2005) 353–360.
- [28] I. Kaplansky, *Commutative rings*, The University of Chicago Press, Chicago, 1974.
- [29] B. G. Kang, *$\star$ -Operations in integral domains*, PhD thesis, The University of Iowa, Iowa City, 1987.
- [30] A. Mimouni, *Integral domains in which each ideal is a w -ideal*, Comm. Algebra **33** (2005) 1345–1355.
- [31] A. Mimouni, *Semistar-operations of finite character on integral domains*, J. Pure Appl. Algebra **200** (2005) 37–50.
- [32] A. Mimouni and M. Samman, *semistar-operations on Valuation domains*, Internat. J. Comm. Rings **2(3)**, (2003) 131–141.
- [33] G. Picozza and F. Tartarone, *When the semistar operation $\bar{*}$ is the identity*, Comm. Algebra to appear.
- [34] M. Zafrullah, *What v -coprimality can do for you, a survey article, to appear in “Multiplicative Ideal Theory: a tribute to the work of Robert Gilmer,”* Eds: J. W. Brewer, S. Glaz, W. J. Heinzer, and B. Olberding, Springer Science+Business Media, 2006.

DEPARTMENT OF MATHEMATICAL SCIENCES, KING FAHD UNIVERSITY OF PETROLEUM & MINERALS, P. O. BOX 5046, DHAHRAN 31261, KSA
E-mail address: `amimouni@kfupm.edu.sa`